

## SPRY2

**Reactivity:** Human Mouse

**Tested applications:** WB

**Recommended Dilution:** WB 1:200 - 1:2000

**Calculated MW:** 35kDa

**Observed MW:** Refer to Figures

**Immunogen:**

A synthetic peptide of human SPRY2

**Storage Buffer:**

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

q

**Synonym:**

hSPRY2;

**Catalog #:** A6262

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 10253

**Isotype:** IgG

**Swiss Prot:** O43597

**Purity:** Affinity purification

For research use only.

**Background:**

This gene encodes a protein belonging to the sprouty family. The encoded protein contains a carboxyl-terminal cysteine-rich domain essential for the inhibitory activity on receptor tyrosine kinase signaling proteins and is required for growth factor stimulated translocation of the protein to membrane ruffles. In primary dermal endothelial cells this gene is transiently upregulated in response to fibroblast growth factor two. This protein is indirectly involved in the non-cell autonomous inhibitory effect on fibroblast growth factor two signaling. The protein interacts with Cas-Br-M (murine) ectropic retroviral transforming sequence, and can function as a bimodal regulator of epidermal growth factor receptor/mitogen-activated protein kinase signaling. This protein may play a role in alveoli branching during lung development as shown by a similar mouse protein.

*To place an order, please [Click HERE](#).*