

## DIDO1

**Reactivity:** Human

**Tested applications:** WB IHC IF

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:20 - 1:50

**Calculated MW:** 243kDa

**Observed MW:** Refer to Figures

**Immunogen:**

Recombinant protein of human DIDO1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

BYE1; DIO1; DATF1; DIDO2; DIDO3; DIO-1; DATF-1; C20orf158; dJ885L7.8;

**Catalog #:** A6453

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 11083

**Isotype:** IgG

**Swiss Prot:** Q9BTC0

**Purity:** Affinity purification

For research use only.

**Background:**

Apoptosis, a major form of cell death, is an efficient mechanism for eliminating unwanted cells and is of central importance for development and homeostasis in metazoan animals. In mice, the death inducer-oblierator-1 gene is upregulated by apoptotic signals and encodes a cytoplasmic protein that translocates to the nucleus upon apoptotic signal activation. When overexpressed, the mouse protein induced apoptosis in cell lines growing in vitro. This gene is similar to the mouse gene and therefore is thought to be involved in apoptosis. Alternatively spliced transcripts have been found for this gene, encoding multiple isoforms.

*To place an order, please [Click HERE](#).*