

## IL17F

**Reactivity:** Human Mouse

**Tested applications:** WB IHC IF

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:100

**Calculated MW:** 18kDa

**Observed MW:** Refer to Figures

**Immunogen:**

Recombinant protein of human IL17F

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

ML1; ML-1; CANDF6; IL-17F;

**Catalog #:** A6487

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 112744

**Isotype:** IgG

**Swiss Prot:** Q96PD4

**Purity:** Affinity purification

For research use only.

**Background:**

The protein encoded by this gene is a cytokine that shares sequence similarity with IL17. This cytokine is expressed by activated T cells, and has been shown to stimulate the production of several other cytokines, including IL6, IL8, and CSF2/GM-CSF. This cytokine is also found to inhibit the angiogenesis of endothelial cells and induce endothelial cells to produce IL2, TGFB1/TGFB, and monocyte chemoattractant protein-1.

*To place an order, please [Click HERE](#).*