

## CCL25

**Reactivity:** Human

**Tested applications:** WB IHC IF

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:100 - 1:200 IF 1:50 - 1:100

**Calculated MW:** 16kDa

**Observed MW:** Refer to Figures

**Immunogen:**

Recombinant protein of human CCL25

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

TECK; Ckb15; SCYA25;

**Catalog #:** A6543

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 6370

**Isotype:** IgG

**Swiss Prot:** O15444

**Purity:** Affinity purification

For research use only.

**Background:**

This antimicrobial gene belongs to the subfamily of small cytokine CC genes. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for dendritic cells, thymocytes, and activated macrophages but is inactive on peripheral blood lymphocytes and neutrophils. The product of this gene binds to chemokine receptor CCR9. Alternative splicing results in multiple transcript variants.

**To place an order, please [Click HERE](#).**