## CTNNBL1

Reactivity: Human Mouse Rat

Tested applications: WB IHC IF

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:65kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human CTNNBL1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Concentration:

b

Synonym:

NAP; P14L; PP8304; C20orf33; dJ633O20.1;

Background:

The protein encoded by this gene is a component of the pre-mRNA-processing factor 19-cell division cycle 5-like (PRP19-CDC5L) protein complex, which activates pre-mRNA splicing and is an integral part of the spliceosome. The encoded protein is also a nuclear localization sequence binding protein, and binds to activation-induced deaminase and is important for antibody diversification. This gene may also be associated with the development of obesity. Alternative splicing results in multiple transcript variants. A pseudogene of this gene has been defined on the X chromosome.

To place an order, please Click HERE.

Polyclonal Antibody

Species:Rabbit

Gene ID:56259

Isotype:IgG

Swiss Prot:Q8WYA6

Purity: Affinity purification

For research use only.





