

## LECT1

**Reactivity:**Human Mouse

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:100

**Calculated MW:**37kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human LECT1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

CHM1; CHM-I; BRICD3; MYETS1;

**Background:**

This gene encodes a glycosylated transmembrane protein that is cleaved to form a mature, secreted protein. The N-terminus of the precursor protein shares characteristics with other surfactant proteins and is sometimes called chondrosurfactant protein although no biological activity has yet been defined for it. The C-terminus of the precursor protein contains a 25 kDa mature protein called leukocyte cell-derived chemotaxin-1 or chondromodulin-1. The mature protein promotes chondrocyte growth and inhibits angiogenesis. This gene is expressed in the avascular zone of prehypertrophic cartilage and its expression decreases during chondrocyte hypertrophy and vascular invasion. The mature protein likely plays a role in endochondral bone development by permitting cartilaginous anlagen to be vascularized and replaced by bone. It may be involved also in the broad control of tissue vascularization during development. Alternative splicing results in multiple transcript variants encoding different isoforms.

**To place an order, please [Click HERE](#).**

**Catalog #:**A6644

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**11061

**Isotype:**IgG

**Swiss Prot:**O75829

**Purity:**Affinity purification

For research use only.