www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

YARS

Reactivity:Human Mouse Rat

Tested applications:WB IHC

 Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

 Calculated MW:59kDa

 Observed MW:Refer to figures

 Immunogen:

 Recombinant protein of human YARS

 Storage Buffer:

 Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

 Synonym:

YRS; YTS; TYRRS; CMTDIC;



Catalog #:A6791 Antibody Type: Polyclonal Antibody Species:Rabbit Gene ID:8565 Isotype:IgG Swiss Prot:P54577 Purity:Affinity purification

For research use only.

Background:

Aminoacyl-tRNA synthetases catalyze the aminoacylation of tRNA by their cognate amino acid. Because of their central role in linking amino acids with nucleotide triplets contained in tRNAs, aminoacyl-tRNA synthetases are thought to be among the first proteins that appeared in evolution. Tyrosyl-tRNA synthetase belongs to the class I tRNA synthetase family. Cytokine activities have also been observed for the human tyrosyl-tRNA synthetase, after it is split into two parts, an N-terminal fragment that harbors the catalytic site and a C-terminal fragment found only in the mammalian enzyme. The N-terminal fragment is an interleukin-8-like cytokine, whereas the released C-terminal fragment is an EMAP II-like cytokine.

To place an order, please Click HERE.



