

## RFC2

---

**Reactivity:**Human Mouse

**Tested applications:**WB

**Recommended Dilution:**WB 1:500 - 1:2000

**Calculated MW:**39kDa

**Observed MW:**Refer to figures

**Immunogen:**

Recombinant protein of human RFC2

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

RFC40;

**Catalog #:**A6847

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**5982

**Isotype:**IgG

**Swiss Prot:**P35250

**Purity:**Affinity purification

For research use only.

**Background:**

This gene encodes a member of the activator 1 small subunits family. The elongation of primed DNA templates by DNA polymerase delta and epsilon requires the action of the accessory proteins, proliferating cell nuclear antigen (PCNA) and replication factor C (RFC). Replication factor C, also called activator 1, is a protein complex consisting of five distinct subunits. This gene encodes the 40 kD subunit, which has been shown to be responsible for binding ATP and may help promote cell survival. Disruption of this gene is associated with Williams syndrome. Alternatively spliced transcript variants encoding distinct isoforms have been described. A pseudogene of this gene has been defined on chromosome 2.

**To place an order, please [Click HERE](#).**