www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

CLCN7

Reactivity: Human Mouse Rat

Tested applications: WB IHC IF

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:100

Calculated MW:89kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human CLCN7

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

CLC7; CLC-7; OPTA2; OPTB4; PPP1R63;

Background:

The product of this gene belongs to the CLC chloride channel family of proteins. Chloride channels play important roles in the plasma membrane and in intracellular organelles. This gene encodes chloride channel 7. Defects in this gene are the cause of osteopetrosis autosomal recessive type 4 (OPTB4), also called infantile malignant osteopetrosis type 2 as well as the cause of autosomal dominant osteopetrosis type 2 (OPTA2), also called autosomal dominant Albers-Schonberg disease or marble disease autosoml dominant. Osteopetrosis is a rare genetic disease characterized by abnormally dense bone, due to defective resorption of immature bone. OPTA2 is the most common form of osteopetrosis, occurring in adolescence or adulthood.

To place an order, please Click HERE.

Antibody Type:

Species: Rabbit

Gene ID:1186

Isotype:IgG

Purity: Affinity purification

For research use only.





