www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

PDHB

Reactivity: Human Mouse Rat

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:39kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human PDHB

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

PDHBD; PHE1B; PDHE1-B;



Catalog #:A6943

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID:5162

Isotype:IgG

Swiss Prot:P11177

Purity: Affinity purification

For research use only.

Background:

The pyruvate dehydrogenase (PDH) complex is a nuclear-encoded mitochondrial multienzyme complex that catalyzes the overall conversion of pyruvate to acetyl-CoA and carbon dioxide, and provides the primary link between glycolysis and the tricarboxylic acid (TCA) cycle. The PDH complex is composed of multiple copies of three enzymatic components: pyruvate dehydrogenase (E1), dihydrolipoamide acetyltransferase (E2) and lipoamide dehydrogenase (E3). The E1 enzyme is a heterotetramer of two alpha and two beta subunits. This gene encodes the E1 beta subunit. Mutations in this gene are associated with pyruvate dehydrogenase E1-beta deficiency. Alternatively spliced transcript variants have been found for this gene.

To place an order, please Click HERE.





