

NAA10

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:100

Calculated MW:26kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human NAA10

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

TE2; ARD1; NATD; ARD1A; ARD1P; DXS707; MCOPS1;

Background:

N-alpha-acetylation is among the most common post-translational protein modifications in eukaryotic cells. This process involves the transfer of an acetyl group from acetyl-coenzyme A to the alpha-amino group on a nascent polypeptide and is essential for normal cell function. This gene encodes an N-terminal acetyltransferase that functions as the catalytic subunit of the major amino-terminal acetyltransferase A complex. Mutations in this gene are the cause of Ogden syndrome. Alternate splicing results in multiple transcript variants.

To place an order, please [Click HERE](#).

Catalog #:A7009

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:8260

Isotype:IgG

Swiss Prot:P41227

Purity:Affinity purification

For research use only.