

## PRC1

---

**Reactivity:**Human

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:**72kDa

**Observed MW:**Refer to figures

**Immunogen:**

Recombinant protein of human PRC1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

1 mg/ml

**Synonym:**

ASE1;

**Catalog #:**A7029

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**9055

**Isotype:**IgG

**Swiss Prot:**O43663

**Purity:**Affinity purification

For research use only.

**Background:**

This gene encodes a protein that is involved in cytokinesis. The protein is present at high levels during the S and G2/M phases of mitosis but its levels drop dramatically when the cell exits mitosis and enters the G1 phase. It is located in the nucleus during interphase, becomes associated with mitotic spindles in a highly dynamic manner during mitosis, and localizes to the cell mid-body during cytokinesis. This protein has been shown to be a substrate of several cyclin-dependent kinases (CDKs). It is necessary for polarizing parallel microtubules and concentrating the factors responsible for contractile ring assembly. Alternative splicing results in multiple transcript variants.

**To place an order, please [Click HERE](#).**