

## NMT2

---

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:100

**Calculated MW:**57kDa

**Observed MW:**Refer to figures

**Immunogen:**

Recombinant protein of human NMT2

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Catalog #:**A7042

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**9397

**Isotype:**IgG

**Swiss Prot:**O60551

**Purity:**Affinity purification

For research use only.

**Background:**

N-myristoyltransferase (NMT) catalyzes the reaction of N-terminal myristoylation of many signaling proteins. It transfers myristic acid from myristoyl coenzyme A to the amino group of a protein's N-terminal glycine residue. Biochemical evidence indicates the presence of several distinct NMTs, varying in apparent molecular weight and /or subcellular distribution. The predicted 498-amino acid of human NMT2 protein shares 77% and 96% sequence identity with human NMT1 and mouse Nmt2 comprise two distinct families of N-myristoyltransferases.

*To place an order, please [Click HERE](#).*