

NCOR1

Reactivity:Human

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:100

Calculated MW:270kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human NCOR1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

N-CoR; TRAC1; N-CoR1; hN-CoR; PPP1R109;

Background:

This gene encodes a protein that mediates ligand-independent transcription repression of thyroid-hormone and retinoic-acid receptors by promoting chromatin condensation and preventing access of the transcription machinery. It is part of a complex which also includes histone deacetylases and transcriptional regulators similar to the yeast protein Sin3p. This gene is located between the Charcot-Marie-Tooth and Smith-Magenis syndrome critical regions on chromosome 17. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 17 and 20.

To place an order, please [Click HERE](#).

Catalog #:A7046

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:9611

Isotype:IgG

Swiss Prot:O75376

Purity:Affinity purification

For research use only.