## CCL26

Reactivity: Human Mouse Rat

Tested applications: WB IHC IF

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:100

Calculated MW:11kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human CCL26

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

IMAC; TSC-1; MIP-4a; SCYA26; MIP-4alpha;

Polyclonal Antibody

Species: Rabbit Gene ID:10344

Isotype:IgG

Swiss Prot:Q9Y258

Purity: Affinity purification

For research use only.

## Background:

This gene is one of two Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 7. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for normal peripheral blood eosinophils and basophils. The product of this gene is one of three related chemokines that specifically activate chemokine receptor CCR3. This chemokine may contribute to the eosinophil accumulation in atopic diseases.

To place an order, please Click HERE.





