OPRM1

Reactivity: Human Mouse Rat

Tested applications:WB

Recommended Dilution: WB 1:500 - 1:1000

Calculated MW:45kDa

Observed MW:Refer to figures

Immunogen:

A synthetic peptide of human OPRM1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

MOP; MOR; LMOR; MOR1; OPRM; M-OR-1;

Polyclonal Antibody Species: Rabbit Gene ID:4988 Isotype:IgG

Swiss Prot:P35372 Purity: Affinity purification

For research use only.

Background:

This gene encodes one of at least three opioid receptors in humans; the mu opioid receptor (MOR). The MOR is the principal target of endogenous opioid peptides and opioid analgesic agents such as beta-endorphin and enkephalins. The MOR also has an important role in dependence to other drugs of abuse, such as nicotine, cocaine, and alcohol via its modulation of the dopamine system. The NM_001008503.2:c.118A>G allele has been associated with opioid and alcohol addiction and variations in pain sensitivity but evidence for it having a causal role is conflicting. Multiple transcript variants encoding different isoforms have been found for this gene. Though the canonical MOR belongs to the superfamily of 7-transmembrane-spanning G-protein-coupled receptors some isoforms of this gene have only 6 transmembrane domains.

To place an order, please Click HERE.



