

BAG2

Reactivity:Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:24kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human BAG2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

BAG-2; dJ41711.2;

Catalog #:A7763

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:9532

Isotype:IgG

Swiss Prot:O95816

Purity:Affinity purification

For research use only.

Background:

BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The predicted BAG2 protein contains 211 amino acids. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner.

To place an order, please [Click HERE](#).