www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

CLTC

Reactivity:Human Mouse Rat

Tested applications:WB IHC

 Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:100

 Calculated MW:192kDa

 Observed MW:Refer to figures

 Immunogen:

 A synthetic peptide of human CLTC

 Storage Buffer:

 Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

 Synonym:

Hc; CHC; CHC17; CLH-17; CLTCL2;

Background:

Clathrin is a major protein component of the cytoplasmic face of intracellular organelles, called coated vesicles and coated pits. These specialized organelles are involved in the intracellular trafficking of receptors and endocytosis of a variety of macromolecules. The basic subunit of the clathrin coat is composed of three heavy chains and three light chains.

To place an order, please Click HERE.



Catalog #:A7886 Antibody Type: Polyclonal Antibody Species:Rabbit Gene ID:1213 Isotype:IgG Swiss Prot:Q00610 Purity:Affinity purification

For research use only.



