www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

BRCC3

Reactivity:Human Mouse Rat

Tested applications:WB IHC

 Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

 Calculated MW:36kDa

 Observed MW:Refer to figures

 Immunogen:

 Recombinant protein of human BRCC3

 Storage Buffer:

 Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

 Synonym:

C6.1A; BRCC36; CXorf53;

Background:

This gene encodes a subunit of the BRCA1-BRCA2-containing complex (BRCC), which is an E3 ubiquitin ligase. This complex plays a role in the DNA damage response, where it is responsible for the stable accumulation of BRCA1 at DNA break sites. The component encoded by this gene can specifically cleave Lys 63-linked polyubiquitin chains, and it regulates the abundance of these polyubiquitin chains in chromatin. The loss of this gene results in abnormal angiogenesis and is associated with syndromic moyamoya, a cerebrovascular angiopathy. Alternative splicing results in multiple transcript variants. A related pseudogene has been identified on chromosome 5.

To place an order, please Click HERE.



Catalog #:A7995 Antibody Type: Polyclonal Antibody Species:Rabbit Gene ID:79184 Isotype:IgG Swiss Prot:P46736 Purity:Affinity purification

For research use only.



