www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

## PDK3

Reactivity:Human Mouse Rat

Tested applications:WB IHC

 Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:100

 Calculated MW:47kDa

 Observed MW:Refer to figures

 Immunogen:

 Recombinant protein of human PDK3

 Storage Buffer:

 Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

 Synonym:

 CMTX6; GS1-358P8.4;



Catalog #:A8028 Antibody Type: Polyclonal Antibody Species:Rabbit Gene ID:5165 Isotype:IgG Swiss Prot:Q15120 Purity:Affinity purification

For research use only.

## Background:

The pyruvate dehydrogenase (PDH) complex is a nuclear-encoded mitochondrial multienzyme complex that catalyzes the overall conversion of pyruvate to acetyl-CoA and CO(2). It provides the primary link between glycolysis and the tricarboxylic acid (TCA) cycle, and thus is one of the major enzymes responsible for the regulation of glucose metabolism. The enzymatic activity of PDH is regulated by a phosphorylation/dephosphorylation cycle, and phosphorylation results in inactivation of PDH. The protein encoded by this gene is one of the three pyruvate dehydrogenase kinases that inhibits the PDH complex by phosphorylation of the E1 alpha subunit. This gene is predominantly expressed in the heart and skeletal muscles. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

To place an order, please Click HERE.



