www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

## PER1

Reactivity:Human Mouse Rat

Tested applications:WB IHC

 Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:100

 Calculated MW:136kDa

 Observed MW:Refer to figures

 Immunogen:

 Recombinant protein of human PER1

 Storage Buffer:

 Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

 Synonym:

PER; hPER; RIGUI;

## Background:

This gene is a member of the Period family of genes and is expressed in a circadian pattern in the suprachiasmatic nucleus, the primary circadian pacemaker in the mammalian brain. Genes in this family encode components of the circadian rhythms of locomotor activity, metabolism, and behavior. This gene is upregulated by CLOCK/ARNTL heterodimers but then represses this upregulation in a feedback loop using PER/CRY heterodimers to interact with CLOCK/ARNTL. Polymorphisms in this gene may increase the risk of getting certain cancers. Alternative splicing has been observed in this gene; however, these variants have not been fully described.

To place an order, please Click HERE.



Catalog #:A8449 Antibody Type: Polyclonal Antibody Species:Rabbit Gene ID:5187 Isotype:IgG Swiss Prot:O15534 Purity:Affinity purification

For research use only.



