

## DDDDK-Tag

---

**Tested applications:**WB IF

**Recommended Dilution:**WB 1:1000 - 1:2000 IF 1:20 - 1:50

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant Protein of human DDDDK

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Background:**

Protein tags are protein or peptide sequences located either on the C- or N- terminal of the target protein, which facilitates one or several of the following characteristics: solubility, detection, purification, localization and expression. The DYKDDDDK(FLAG) peptide has been used extensively as a general tag in expression vectors. This peptide can be expressed and detected with the protein of interest as an amino-terminal or carboxy-terminal fusion. N-terminal FLAG vectors provide an Ek cleavage site for removal of the fusion tag. The FLAG peptide is likely to be located on the surface of a fusion protein because of its hydrophilic nature. As a result, the FLAG peptide is more likely to be accessible to antibodies. A DDDDK can be used in many different assays that require recognition by an antibody, such as western blotting, immunocytochemistry, immunoprecipitation, flow cytometry, protein purification, and in the study of protein-protein interactions, cell ultrastructure, and protein localization and so on. This antibody is a mouse monoclonal antibody raised against 3xFlag (3xDYKDDDDKT) sequence and recognizes the (3x)DYKDDDDK peptide and detects DDDDKged proteins.

**To place an order, please [Click HERE](#).**

**Catalog #:**AE004

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Isotype:**IgG

**Purity:**Affinity purification

For research use only.