Phospho-LCK-Y505

Reactivity: Human Mouse

Tested applications:WB

Recommended Dilution: WB 1:500 - 1:2000

Calculated MW:56kDa

Observed MW:Refer to Figures

Immunogen:

A phospho specific peptide corresponding to residues surrounding Y505 of human LCK

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Concentration:

bd

Synonym:

LSK; YT16; p56lck; pp58lck;

Background:

This gene is a member of the Src family of protein tyrosine kinases (PTKs). The encoded protein is a key signaling molecule in the selection and maturation of developing T-cells. It contains N-terminal sites for myristylation and palmitylation, a PTK domain, and SH2 and SH3 domains which are involved in mediating protein-protein interactions with phosphotyrosine-containing and proline-rich motifs, respectively. The protein localizes to the plasma membrane and pericentrosomal vesicles, and binds to cell surface receptors, including CD4 and CD8, and other signaling molecules. Multiple alternatively spliced variants, encoding the same protein, have been described.

To place an order, please Click HERE.



Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID:3932

Isotype:IgG

Swiss Prot:P06239

Purity: Affinity purification

For research use only.





