

Phospho-MDM2-S166

Reactivity:Human

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:100 IF 1:100 - 1:200

Calculated MW:90kDa

Observed MW:Refer to Figures

Immunogen:

A phospho specific peptide corresponding to residues surrounding S166 of human MDM2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

fl

Synonym:

HDMX; hdm2; ACTFS;

Catalog #:AP0293

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:4193

Isotype:IgG

Swiss Prot:Q00987

Purity:Affinity purification

For research use only.

Background:

This gene encodes a nuclear-localized E3 ubiquitin ligase. The encoded protein can promote tumor formation by targeting tumor suppressor proteins, such as p53, for proteasomal degradation. This gene is itself transcriptionally-regulated by p53. Overexpression or amplification of this locus is detected in a variety of different cancers. There is a pseudogene for this gene on chromosome 2. Alternative splicing results in a multitude of transcript variants, many of which may be expressed only in tumor cells.

To place an order, please [Click HERE](#).