

## Phospho-CDKN1A-T145

**Reactivity:** Human

**Tested applications:** IHC

**Recommended Dilution:** IHC 1:50 - 1:100

**Calculated MW:** 21kDa

**Observed MW:** Refer to Figures

**Immunogen:**

A phospho specific peptide corresponding to residues surrounding T145 of human CDKN1A

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

b

**Synonym:**

P21; CIP1; SDI1; WAF1; CAP20; CDKN1; MDA-6; p21CIP1;

**Catalog #:** AP0327

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 1026

**Isotype:** IgG

**Swiss Prot:** P38936

**Purity:** Affinity purification

For research use only.

**Background:**

This gene encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-CDK2 or -CDK4 complexes, and thus functions as a regulator of cell cycle progression at G1. The expression of this gene is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen (PCNA), a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of CDK2, and may be instrumental in the execution of apoptosis following caspase activation. Multiple alternatively spliced variants have been found for this gene.

**To place an order, please [Click HERE](#).**