

## Borrelia OspC

**Description:** Recombinant *Borrelia burgdorferi* Outer Surface Protein C produced in *E. coli* is a non-glycosylated, polypeptide chain having a calculated molecular mass of 21,443 Dalton. *Borrelia* OspC is expressed with a -6x His tag at N-terminus and purified by proprietary chromatographic techniques.

Catalog #: BOPS-011

For research use only.

**Source:** *Escherichia Coli*.

**Physical Appearance:** Sterile Filtered clear solution.

**Purity:** Greater than 80.0% as determined by SDS-PAGE.

**Formulation:**

*Borrelia* OspC (1.74mg/1ml) is supplied in 16mM HEPES buffer pH-7.0, 300mM NaCl and 20% glycerol.

**Stability:**

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. Avoid multiple freeze-thaw cycles.

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Applications:**

Western blot with Lyme positive plasma.

**Introduction:**

*Borrelia* belongs to a genus of bacteria of the spirochete phylum. *Borrelia* causes borreliosis, which is a zoonotic, vector-borne disease transmitted mainly by ticks and some by lice, depending on the species. Of the 36 known species of *Borrelia*, 12 are distinguished to cause Lyme disease or borreliosis and are transmitted by ticks. The main *Borrelia* species causing Lyme disease are *Borrelia burgdorferi*, *Borrelia afzelii*, and *Borrelia garinii*. The *Borrelia* genus members have a linear chromosome which is about 900 kbp in length as well as an excess of both linear and circular plasmids in the 5-220 kbp size range. The plasmids are atypical, as compared to most bacterial plasmids, since they contain many paralogous sequences, a large number of pseudogenes and, in some cases, essential genes. Moreover, a number of the plasmids have features suggesting that they are prophages.

**To place an order, please [Click HERE](#).**