

## CCL3L1 Human

**Description:**CCL3L1 Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 70 amino acids and having a molecular mass of 7.7kDa. The CCL3L1 is purified by proprietary chromatographic techniques.

**Catalog #:**CHPS-270

**Synonyms:**C-C motif chemokine 3-like 1, G0/G1 switch regulatory protein 19-2, LD78-beta(1-70), PAT 464.2, Small-inducible cytokine A3-like 1, Tonsillar lymphocyte LD78 beta protein, CCL3L1, D17S1718, G0S19-2, SCYA3L1, LD78, 464.2, SCYA3L, LD78BETA, MGC12815.

For research use only.

**Source:**Escherichia Coli.

**Physical Appearance:**Sterile Filtered White lyophilized (freeze-dried) powder.

**Amino Acid Sequence:**APLAADTPTA CCFSYTSRQI PQNFIADYFE TSSQCSKPSV  
IFLTRGRQV CADPSEEWVQ KYVSDLELSA.

**Purity:**Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

**Formulation:**

Lyophilized from a 0.2

**Stability:**

Lyophilized CCL3L1 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CCL3L1 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Solubility:**

It is recommended to reconstitute the lyophilized CCL3L1 in sterile 18M-cm H2O not less than 100

**Introduction:**

CCL3L1 is a small cytokine that belongs to the CC chemokines. The CCL3L1 gene is one of several cytokine genes clustered on the q-arm of chromosome 17. CCL3L1 is involved in immunoregulatory and inflammatory processes. CCL3L1 binds to several chemokine receptors including CCBP2 and CCR5. CCR5 is a co-receptor for HIV, and binding of the CCL3L1 protein to CCR5 inhibits HIV entry.

**Biological Activity:**

The biological activity was determined by its ability to chemoattract human monocytes using a concentration range of 1.0-10.0 ng/ml.

**To place an order, please [Click HERE](#).**