

IL 3 Human, HEK

Description: Interleukin-3 Human Recombinant produced in HEK cells is a glycosylated monomer, having a molecular weight range of 17-45kDa due to glycosylation. The IL-3 is purified by proprietary chromatographic techniques.

Catalog #: CYPs-103

For research use only.

Synonyms: MCGF (Mast cell growth factor), Multi-CSF, HCGF, P-cell stimulation factor, IL-3, MGC79398, MGC79399.

Source: HEK.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Purity: Greater than 90% as observed by SDS-PAGE.

Formulation:

The IL3 was lyophilized from 1mg/ml in 1xPBS.

Stability:

Lyophilized IL-3 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL3 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolabs products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized IL-3 in sterile water not less than 100

Introduction:

IL3 is a potent growth promoting cytokine. This cytokine is capable of supporting the proliferation of a broad range of hematopoietic cell types. It is involved in a variety of cell activities such as cell growth, differentiation and apoptosis. This cytokine has been shown to also possess neurotrophic activity, and it may be associated with neurologic disorders.

Biological Activity:

The activity was determined by the dose dependent stimulation of the proliferation of human TF-1 cells (human erythroleukemic indicator cell line) and is typically 0.5-2ng/ml corresponding to a specific activity of 500,000-2,000,000units/mg.

To place an order, please [Click HERE](#).