

IL 10 Human, HEK

Description: Interleukin-10 Human Recombinant produced in HEK cells is a glycosylated non-disulfide bonded homodimer, having a total molecular weight of 17kDa. The IL10 is purified by proprietary chromatographic techniques.

Catalog #: CYPs-107

Synonyms: B-TCGF, CSIF, TGIF, IL-10, IL10A, MGC126450, MGC126451, Cytokine synthesis inhibitory factor.

For research use only.

Source: HEK.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Purity: Greater than 95% as observed by SDS-PAGE.

Formulation:

The IL10 was lyophilized from 1mg/ml in 10mM Tris-HCl, pH 7.4 and 150mM NaCl.

Stability:

Lyophilized IL-10 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL10 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolabs products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized IL-10 in sterile water not less than 100

Introduction:

IL10 is a cytokine produced primarily by monocytes and to a lesser extent by lymphocytes. This cytokine has pleiotropic effects in immunoregulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II Ags, and costimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. This cytokine can block NF-kappa B activity, and is involved in the regulation of the JAK-STAT signaling pathway. Knockout studies in mice suggested the function of this cytokine as an essential immunoregulator in the intestinal tract.

Biological Activity:

The specific activity was determined by the dose-dependent stimulation of the proliferation of MC/9 cells (mouse mast cell line) in the presence of 200 pg/ml IL-4, and is typically 0.3-1.5ng/ml.

To place an order, please [Click HERE](#).