

IL36A 158 a.a. Human

Description:IL36A 158 a.a. Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 158 amino acids and having a molecular mass of 17.7kDa. The IL36A 158 a.a. Human is purified by proprietary chromatographic techniques.

Synonyms:Interleukin 36 alpha, FIL1E, IL1F6, FIL1, IL1(EPSILON), interleukin 1 family member 6 (epsilon), MGC129552, MGC129553.

Source:Escherichia Coli.

Physical Appearance:Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence:MEKALKIDTP QQGSIQDINH RVWVLQDQTL IAVPRKDRMS
PVTIALISCR HVETLEKDRG NPIYLG LNL NLCLMCAKVG DQPTLQLKEK DIMDLYNQPE
PVKSFLFYHS QSGRNSTFES VAFPGWFIAS SSEGGCPLIL TQELGKANTT DFGLTMLF

Purity:Greater than 95.0% as determined by SDS-PAGE and HPLC analyses.

Formulation:

Lyophilized from a 0.2m filtered concentrated solution in 2xPBS, pH 7.4.

Stability:

Lyophilized IL36A 158 a.a. Human although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL36A 158 a.a. Human should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized IL36A 158 a.a. Human in sterile 18M-cm H2O not less than 100

Introduction:

Human IL-36a belongs to the IL-1 family which includes IL-1b, IL-1a, IL-1ra, IL-18, IL-36ra (IL1F5), IL-36b (IL1F8), IL-36g (IL1F9), IL-37 (IL1F7) and IL-38 (IL-1F10). The IL-1 family members display a 12 b-strand, b-trefoil configuration, and are thought to have ascended from a mutual ancestral gene. IL-36a is an 18-22kDa, 158aa intracellular and secreted protein which holds no signal sequence, no prosegment and no potential from N-linked glycosylation sites. IL-36a is released as a reaction to LPS and the cell ATP-induced activation of the P2X7 receptor. Human IL-36a (aa 6-158) shares 57-68% aa sequence homology with mouse, rabbit, equine and bovine IL-36a and 27-57% aa sequence homology with other new IL-1 family members. IL-36a is mostly found in skin and lymphoid tissues, but also in fetal brain, trachea, stomach and intestine.

Biological Activity:

Fully biologically active when compared to standard. The specific activity determined by its ability in a functional ELISA. Immobilized rHuIL-36 at 1 g/mL can bind recombinant human IL-1 Rrp2 Fc Chimera with a range of 0.15-5g/ml corresponding to a specific activity of

www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom
200,000-6,666,667IU/mg.



To place an order, please [Click HERE](#).

Catalog #:CYP5-186

For research use only.

© 2015 NeoScientific



neobiolab | [Term](#) | [Quote/Order](#) | [Services](#) | [Products](#) | [Support](#) | [Corporate](#) | [Contact Us](#)



Copyrights @ 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)

Request a
CATALOG
[CLICK HERE](#)