www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

IL 7 Human, HEK

Description: IL-7 Human Recombinant produced in HEK cells is a glycosylated monomer, having a molecular weight range of 19-30kDa due to glycosylation. The IL-7 is purified by proprietary chromatographic techniques.

Catalog #:CYPS-204

For research use only.

Synonyms:Lymphopoietin 1 (LP-1), pre-B cell factor, IL-7.

Source: HEK.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Purity: Greater than 95% as obsereved by SDS-PAGE.

Formulation:

The IL7 was lyophilized from 1mg/ml in 1xPBS.

Stability:

Lyophilized IL-7 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL7 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolabs products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized IL-7 in sterile PBS containing 0.1% endotoxin-free recombinant HSA.

Introduction:

IL-7 is a cytokine important for B and T cell development. This cytokine and the hepatocyte growth factor (HGF) form a heterodimer that functions as a pre-pro-B cell growth-stimulating factor. This cytokine is found to be a cofactor for V(D)J rearrangement of the T cell receptor beta (TCRB) during early T cell development. This cytokine can be produced locally by intestinal epithelial and epithelial goblet cells, and may serve as a regulatory factor for intestinal mucosal lymphocytes. Knockout studies in mice sµggested that this cytokine plays an essential role in lymphoid cell survival.

Biological Activity:

The activity was determined by the dose dependent stimulation of the proliferation of murine 2E8 cells and is typically 0.6ng/ml corresponding to a specific activity of 1,666,667units/mg.

To place an order, please Click HERE.





