

IL 3 Mouse

Description: Interleukin-3 mouse Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 135 amino acids and having a molecular mass of 15100 Dalton. The IL-3 is purified by proprietary chromatographic techniques.

Synonyms: MCGF (Mast cell growth factor), Multi-CSF, HCGF, P-cell stimulation factor, IL-3, MGC79398, MGC79399.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: The sequence of the first five N-terminal amino acids was determined and was found to be Met-Asp-Thr-His-Arg.

Purity: Greater than 97.0% as determined by: (a) Analysis by SEC-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

Lyophilized from a concentrated (1mg/ml) solution in water containing no additives.

Stability:

Lyophilized Interleukin-3 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL-3 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Interleukin-3 in sterile 18M-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Introduction:

Interleukin-3 is a pleiotropic cytokine produced primarily by activated T cells. IL-3 is thought to function via specific cell surface receptors to stimulate the proliferation, differentiation and survival of haematopoietic cell lines. IL-3 has also been shown to affect the functional activity of a variety of other cell types including mast cells, eosinophils, megakaryocytes and basophils.

Biological Activity:

The ED₅₀ as determined by the dose-dependant stimulation of murine M-NFS-60 cells is < 0.05 ng/ml, corresponding to a Specific Activity of 20,000,000IU/mg.

References:

1. Title: Direct binding of Cbl to Y568 and Y936 of the stem cell factor receptor/c-Kit is required for ligand-induced ubiquitination, internalization and degradation. Publication: Biochemical Journal Immediate Publication. Published on 19 Jun 2006 as manuscript

www.neobiolab.com

info@neobiolab.com

888.754.5670, +1 617.500.7103 United States

0800.088.5164, +44 020.8123.1558 United Kingdom

BJ20060464.Link:http://www.biochemj.org/bj/imps_x/pdf/BJ20060464.pdf2.Title:enhanced self-renewal of hematopoietic stem/progenitor cells mediated by the stem cell gene

sall4.Publication:Link:<http://www.biomedcentral.com/content/pdf/1756-8722-4-38.pdf>3.Title:Identification of Y589 and Y599 in the juxtamembrane domain of Flt3 as ligand-induced

autophosphorylation sites involved in binding of Src family kinases and the protein tyrosine

phosphatase SHP2.Publication:Published online before printMay 9, 2006,

doi:10.1182/blood-2005-07-008896Submitted: July 14, 2005Accepted: April 26, 2006

BloodSeptember 1, 2006vol. 108no.

51542-1550.Link:<http://bloodjournal.hematologylibrary.org/content/108/5/1542.full>



Catalog #:CYPs-378

For research use only.

To place an order, please [Click HERE](#).

© 2015 NeoScientific



neobiolab | Term | Quote/Order | Services | Products | Support | Corporate | Contact Us

Copyrights @ 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)



Request a
CATALOG
[CLICK HERE](#)