

IL 1 alpha Porcine

Description: Interleukin-1A Porcine Recombinant produced in E.Coli is single, a non-glycosylated, Polypeptide chain containing 158 amino acids and having a molecular mass of 18076 Dalton. The IL-1A is purified by proprietary chromatographic techniques.

Catalog #: CYPs-403

For research use only.

Synonyms: Hematopoietin-1, Lymphocyte-activating factor (LAF), Endogenous Pyrogen (EP), Leukocyte Endogenous Mediator (LEM), Mononuclear Cell Factor (MCF), IL-1 alpha, IL1, IL-1A, IL1F1.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: The sequence of the first five N-terminal amino acids was determined and was found to be Ser-Ala-Thr-Tyr-Ser.

Purity: Greater than 95.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

The protein was lyophilized from a concentrated (1mg/ml) sterile solution containing no additives.

Stability:

Lyophilized Interleukin-1 alpha although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL1A should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Interleukin-1 alpha in sterile 18M-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Introduction:

Interleukin-1 alpha is a proinflammatory cytokine produced by a wide variety of cell types, including macrophages, osteoblasts, monocytes and hepatocytes. Circulating levels of are normally low and only rise after stimulation by agents such as those produced by inflammation, infection or microbial endotoxins. IL-1 alpha possesses a wide variety of biological activities and exerts its effects by binding to specific cell surface receptors.

Biological Activity:

The ED₅₀ as determined by the dose-dependant stimulation of D10S cells is < 0.03 ng/ml.

To place an order, please [Click HERE](#).