

IL 4 Human, His

Description: Interleukin-4 Human Recombinant produced in E.Coli is single, a non-glycosylated, Polypeptide chain containing 150 amino acids fragment (25-153) and having a total molecular mass of 17.2kDa. The IL-4 is fused to a 20 amino acid His-Tag at N-terminus and purified by proprietary chromatographic techniques.

Catalog #: CYP5-490

For research use only.

Synonyms: BCGF, BCDF, B cell stimulating factor, BSF-1, Lymphocyte stimulatory factor 1, IL-4, MGC79402, Binetrakin, Pitrakinra.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered clear solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MHKCDITLQE IIKTLNSLTE
QKTLCTELTV TDIFAASKNT TEKETFCAA TVLRQFYSHH EKDTRCLGAT AQQFHRHKQL
IRFLKRLDRN LWGLAGLNSC PVKEANQSTL ENFLERLKI MREKYSKCSS.

Purity: Greater than 95.0% as determined by SDS-PAGE.

Formulation:

Interleukin-4 His-Tag is supplied in 20mM Tris-HCl and 10% glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please avoid freeze thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Interleukin-4 is a pleiotropic cytokine produced primarily by activated T lymphocytes, basophils and mast cells. Multiple immune response-modulating functions are performed by IL-4 on a variety of cell types and it has an important role in the regulator of isotype switching, induction of IgE production in B lymphocytes and differentiation of precursor T helper cells. IL-4 binds to both membrane-bound and secreted soluble IL-4 receptors.

Biological Activity:

The ED50 for this effect is <1.8ng/ml. Measured in a cell proliferation assay using TF1 human erythroleukemic cells.

To place an order, please [Click HERE](#).