

SHH Mouse

Description: Sonic Hedgehog Recombinant Mouse produced in E.Coli is a single, non-glycosylated polypeptide chain containing 176 amino acids and having a molecular mass of 19.8 kDa. The Mouse Sonic Hedgehog is 99% homologous to the human gene. Cysteine at position 25 has been substituted with Ile. The Sonic Hedgehog is purified by proprietary chromatographic techniques.

Synonyms: SHH, HHG-1, HHG1, Sonic hedgehog protein.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: MIIGPGRGFG KRRHPKKLTP LAYKQFIPNV AEKTLGASGR
YEGKITRNS RFKELTPNYN PDIIKFDEEN TGADRLMTQR CKDKLNALAI SVMNQWPGVK
LRVTEGWDED GHHSSESLHY EGRAVDITTS DRDRSKYGML ARLAVEAGFD WVYYESKAHI
HCSVKAENSV AAKSGG.

Purity: Greater than 97.0% as determined by SDS-PAGE.

Formulation:

Lyophilized from a concentrated (1mg/ml) solution in water containing 10mM Na₂PO₄ pH-7.5.

Stability:

Lyophilized Mouse Sonic Hedgehog although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Sonic Hedgehog should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Sonic Hedgehog in sterile 18M-cm H₂O not less than 100

Introduction:

Recombinant Mouse Sonic Hedgehog is part of a small group of secreted proteins that are vital for development in both vertebrates and invertebrates. 3 mammalian hedgehog genes (sonic, desert, Indian) share about 60% homology. The Mouse Sonic Hedgehog is 99% homologous to the human gene. Sonic Hedgehog is a protein that is vital in guiding the early embryo. It has been associated as the major inductive signal in patterning of the ventral neural tube, the anterior-posterior limb axis, and the ventral somites. Sonic Hedgehog binds to the patched receptor, which functions in association with smoothened, to activate the transcription of target genes. In the absence of sonic Hedgehog, patched receptor represses the constitutive signaling activity of smoothened. Sonic Hedgehog also regulates another factor, the gli oncogene. Sonic Hedgehog intercellular signal is essential for a various patterning events during development: signal produced by the notochord that induces ventral cell fate in the neural tube and somites, and the polarizing signal for patterning of the anterior-posterior axis of the developing limb bud. Sonic

www.neobiolab.com

info@neobiolab.com

888.754.5670, +1 617.500.7103 United States

0800.088.5164, +44 020.8123.1558 United Kingdom

HedgeHog exhibits both floor plate- and motor neuron-inducing activity. Mutations in a long-range Sonic HedgeHog enhancer located in an intron of the limb region 1 gene result in preaxial polydactyly.



Catalog #:CYP5-604

Biological Activity:

Determined by the dose-dependent induction of alkaline phosphatase production by C3H/10T1/2 (CCL-226) fibroblasts and is typically 0.48-0.72

For research use only.

To place an order, please [Click HERE](#).