

IL 3 Dog

Description: Interleukin-3 Canine Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 120 amino acids and having a molecular mass of 14 kDa. The IL-3 is purified by proprietary chromatographic techniques.

Catalog #: CYP5-729

Synonyms: MCGF (Mast cell growth factor), Multi-CSF, HCGF, P-cell stimulation factor, IL-3, MGC79398, MGC79399.

For research use only.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: RPFSTDLPKQ YFTMINEIME MLNKSPSPSE EPLDSNEKET
LLEDTLRPN LDVFLNASSK FHKNGLLIWN NLKEFLPLLP TPTPRGEPIS IMENNWGDFQ
RKLLKYLEAL DNFLNFKNKP.

Purity: Greater than 97.0% as determined by: (a) Analysis by HPLC. (b) Analysis by SDS-PAGE.

Formulation:

Lyophilized from a concentrated (1mg/ml) solution in water containing 1X PBS pH7.4.

Stability:

Lyophilized IL3 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL3 should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized IL3 in sterile 18M-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Introduction:

IL3 is a potent growth promoting cytokine. This cytokine is capable of supporting the proliferation of a broad range of hematopoietic cell types. It is involved in a variety of cell activities such as cell growth, differentiation and apoptosis. This cytokine has been shown to also possess neurotrophic activity, and it may be associated with neurologic disorders.

Biological Activity:

The ED50 as determined by the dose-dependant stimulation of TF-1 cells is
To place an order, please [Click HERE](#).