

## HSD17B14 Human

**Description:** HSD17B14 Human Recombinant fused with a 36 amino acid His tag at N-terminus produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 306 amino acids (1-270 a.a.) and having a molecular mass of 32.4kDa. The HSD17B14 is purified by proprietary chromatographic techniques.

**Catalog #:** ENPS-036

For research use only.

**Synonyms:** 17-beta-hydroxysteroid dehydrogenase 14, 17-beta-HSD 14, 17-beta-hydroxysteroid dehydrogenase DHRS10, Dehydrogenase/reductase SDR family member 10, Retinal short-chain dehydrogenase/reductase retSDR3, HSD17B14, DHRS10, SDR3, SDR47C1, retSDR3.

**Source:** Escherichia Coli.

**Physical Appearance:** Sterile Filtered colorless solution.

**Amino Acid Sequence:** MRGSHHHHHH GMASMTGGQQ MGRDLYDDDD KDRWGSMATG  
TRYAGKVVVV TGGGRGIGAG IVRAFVNSGA RVVICDKDES GGRALEQELP GAVFILCDVT  
QEDDVKTLS ETIRRFGRLD CVVNNAGHHP PPQRPEETSA QGFRQLLELN LLGTYTLTKL  
ALPYLRKSQG NVINISLVG AIGQAQAVPY VATKGAVTAM TKALALDESP YGVRVNCISP  
GNIWTPLWEE LA

**Purity:** Greater than 95.0% as determined by SDS-PAGE.

### Formulation:

The HSD17B14 solution (1 mg/ml) contains 20mM Tris-HCl buffer (pH8.0), 20% glycerol, 0.1M NaCl and 1mM DTT.

### Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

### Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

### Introduction:

17-beta-hydroxysteroid dehydrogenase 14 (HSD17B14) is a member of the 17-beta-HSD family of proteins, which regulate the availability of steroids within various tissues throughout the body. 17-beta-hydroxysteroid dehydrogenases (HSD17B14) are mainly involved in metabolism of steroids at the C17 position and also of other substrates, such as fatty acids, prostaglandins, and xenobiotics. HSD17B14 exists as a homotetramer that localizes to the cytoplasm and is highly expressed in the brain, placenta, liver and kidney.

**To place an order, please [Click HERE](#).**