

HDHD2 Human

Description:HDHD2 produced in E.Coli is a single, non-glycosylated polypeptide chain containing 279 amino acids (1-259a.a.) and having a molecular mass of 30.6kDa.HDHD2 is fused to a 20 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #:ENPS-045

For research use only.

Synonyms:Haloacid Dehalogenase-like Hydrolase Domain Containing 2, DKFZP564D1378, 3110052N05Rik, DKFZp564D1378.

Source:Escherichia Coli.

Physical Appearance:Sterile Filtered clear solution.

Amino Acid Sequence:MGSSHHHHHH SSSLVPRGSH MAACRALKAV LVDL SGLHI
EAAVPGAQE ALKRLRGASV IIRFVTNTTK ESKQDLLERL RKLEFDISED EIFTSLTAAR
SLLERKQVRP MLLVDDRALP DFKGIQTSDP NAVVMGLAPE HFHYQLNQA FRLLLDGAPL
IAIHKARYYK RKGDLALGPG PFVTALEYAT DTKATVVGKP EKTFLEALR GTGCEPEEAV
MIGDDCRDDV GG

Purity:Greater than 95% as determined by SDS-PAGE.

Formulation:

The HDHD2 protein solution (1mg/1ml) is formulated in 20mM Tris-HCl buffer (pH 8.0), 0.1M NaCl, 1mM DTT, and 10% glycerol.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

HDHD2 is a member of the HAD-like hydrolase superfamily which includes L-2-haloacid dehalogenase, epoxide hydrolases and phosphatases. There are two active sites in HDHD2 - an L-2-haloacid dehalogenase and a carboxylate group. The L-2-haloacid dehalogenase active site catalyzes the hydrolytic dehalogenation of D- and L-2-haloalkanoic acids, producing L- and D-2-hydroxyalkanoic acids.

Storage:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time.Please avoid freeze thaw cycles.

To place an order, please [Click HERE](#).