www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

## ATP50 Human

Description: ATP5O Human Recombinant fused with a 21 amino acid His tag at N-terminus produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 211 amino acids (24-213 a.a.) and having a molecular mass of 23.1kDa. The ATP5O is purified by proprietary chromatographic techniques.

Catalog #:ENPS-050

For research use only.

Synonyms: ATP synthase subunit O mitochondrial, Oligomycin sensitivity conferral protein, OSCP, ATP50, ATPO.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MFAKLVRPPV QVYGIEGRYA TALYSAASKQ NKLEQVEKEL LRVAQILKEP KVAASVLNPY VKRSIKVKSL NDITAKERFS PLTTNLINLL AENGRLSNTQ GVVSAFSTMM SVHRGEVPCT VTSASPLEEA TLSELKTVLK SFLSQGQVLK LEAKTDPSIL GGMIVRIGEK YVDMSVKTKI QKLGRAMREI V.

Purity: Greater than 95.0% as determined by SDS-PAGE.

### Formulation:

The ATP5O solution (1 mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 1mM DTT, 40% glycerol and 0.2M NaCl.

# Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

## Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

## Introduction:

ATP synthase subunit O (ATP50) localizes to the mitochondria and catalyzes ATP synthesis. ATP50 is a component of the F-type ATPase found in the mitochondrial matrix. F-type ATPases are composed of a catalytic core and a membrane proton channel. ATP5O seems to be part of the connector connecting these two components and may be involved in transmission of conformational changes or proton conductance.

To place an order, please Click HERE.





