www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom

AKR1D1 Human

Catalog #:ENPS-105

For research use only.

Description:AKR1D1 Human Recombinant fused with a 20 amino acid His tag at N-terminus produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 346 amino acids (1-326 a.a.) and having a molecular mass of 39.5kDa. The AKR1D1 is purified by proprietary chromatographic techniques.

Synonyms:3-oxo-5-beta-steroid 4-dehydrogenase, Aldo-keto reductase family 1 member D1, Delta(4)-3-ketosteroid 5-beta-reductase, Delta(4)-3-oxosteroid 5-beta-reductase, KR1D1, SRD5B1, CBAS2, 3o5bred.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence:MGSSHHHHHH SSGLVPRGSH MDLSAASHRI PLSDGNSIPI IGLGTYSEPK STPKGACATS VKVAIDTGYR HIDGAYIYQN EHEVGEAIRE KIAEGKVRRE DIFYCGKLWA TNHVPEMVRP TLERTLRVLQ LDYVDLYIIE VPMAFKPGDE IYPRDENGKW LYHKSNLCAT WEAMEACKDA GLVKSLGVSN FNRRQLELIL NKPGLKHKPV SNQVECHPYF TQPKLLKFCQ QH

Purity: Greater than 90.0% as determined by SDS-PAGE.

Formulation:
The AKR1D1 solution (0.5 mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 1mM DTT, 20% glycerol and 100mM NaCl.

Stability:
Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles.

Usage:
NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drμgs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:
Aldo-keto reductase family 1 member D1 (AKR1D1) belongs to the AKR superfamily. The AKR family proteins are soluble NADPH oxidoreductases, which have vital roles in the metabolism of drμgs, carcinogens and reactive aldehydes. AKR1D1 is also responsible for the catalysis of the 5-beta-reduction of bile acid intermediates and steroid hormones that carry a delta (4)-3-1 structure. AKR1D1 is highly expressed in the liver, colon and testis. Deficiency of the AKR1D1 enzyme may contribute to hepatic dysfunction.

To place an order, please Click HERE.

© 2015 NeoScientific

Copyrights @ 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)
Request a CATALOG CLICK HERE