

MMP-2 Human

Description:MMP-2 Human Recombinant produced in HEK293 cells is a proform of the Human MMP-2 (Ala30-Cys660) and fused with a ployhistide tag at the C-terminus, having an Mw of 71kDa. MMP-2 is purified by proprietary chromatographic techniques.

Catalog #:ENPS-107

Synonyms:72 kDa type IV collagenase, 72 kDa gelatinase, Gelatinase A, Matrix metalloproteinase-2, MMP-2, TBE-1, MMP2, CLG4A, CLG4, MONA, MMP-II.

For research use only.

Source:HEK293 cells.

Physical Appearance:The MMP-2 is supplied as a sterile Filtered colorless solution.

Purity:Greater than 95% as determined by SDS-PAGE.

Formulation:

The MMP-2 is supplied as a 0.2

Stability:

Store MMP-2 at 4°C if entire vial will be used within 2-4 weeks. Store frozen at -20°C for longer periods of time. Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Matrix metalloproteinase-2 (MMP-2) is a type IV collagenase, which is involved in endometrial menstrual breakdown, regulation of vascularization and the inflammatory response. MMP-2 contains a number of distinct domains: a prodomain that is cleaved upon activation; a catalytic domain containing the zinc binding site; a fibronectin like domain believed to have a role in substrate targeting; and a carboxyl terminal (hemopexin like) domain containing 2 N-linked glycosylation. The MMP-2 can degrade an extensive array of substrates including type IV, V, VII and X collagens as well as gelatin type I. In addition, MMP-2 interacts with THBS2, TIMP2, Thrombospondin 1, CCL7 and TIMP4. MMP-2 autocatalytic cleavage in the C-terminal generates the anti-angiogenic peptide, PEX. This process seems to be made possible by binding integrin α v/ β 3. Defects in the MMP-2 are the cause of Torg-Winchester syndrome (TWS), aka multicentric osteolysis nodulosis and arthropathy (MONA).

Biological Activity:

The activity was measured by its ability to cleave the colorimetric peptide substrate, Mca-PLGL-DpaAR-NH₂, The specific activity is > 1,000 pmoles/min/

References:

Title:MMP-2 regulates human platelet activation by interacting with integrin α IIb β 3. Publication: To cite this article: Choi W-S, Jeon O-H, Kim H-H, Kim D-S. MMP-2 regulates human platelet activation by interacting with integrin α IIb β 3. J Thromb Haemost 2008; 6: 51723. Link: <http://onlinelibrary.wiley.com/doi/10.1111/j.1538-7836.2007.02871.x/pdf>

To place an order, please [Click HERE](#).