

IDI2 Human

Description:IDI2 produced in E.Coli is a single, non-glycosylated polypeptide chain containing 247 amino acids (1-227 a.a.) and having a molecular mass of 28.9kDa. IDI2 is fused to a 20 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #:ENPS-114

Synonyms:Isopentenyl-diphosphate Delta-isomerase 2, Isopentenyl pyrophosphate isomerase 2, IPP isomerase 2, IPPI2, IDI2.

For research use only.

Source:Escherichia Coli.

Physical Appearance:Sterile filtered colorless solution.

Amino Acid Sequence:MGSSHHHHH SSGLVPRGSH MSDINLDWVD RRQLQRLEEM
LIVVDENDKV IGADTKRNCH LNIENIEKGLL HRAFSVVLFN TKNRILIQQR SDTKVTFPGY
FTDSCSSHPL YNPAELEEKD AIGVRRAAQR RLQAELGIPG EQISPEDIVF MTIYHHKAKS
DRIWGEHEIC YLLLVRKNVT LNPDPSETKS ILYLSQEELW ELLEREARGE VKVTPWLRTI
AERFLYRWWP HL

Purity:Greater than 80.0% as determined by SDS-PAGE.

Formulation:

IDI2 solution (0.5mg/ml) containing 20mM Tris-HCl buffer (pH8.0), 10% glycerol, 1mM DTT and 0.1mM PMSF.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. They may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Isopentenyl-diphosphate Delta-isomerase 2 (IDI2) is a member of the IPP isomerase type 1 family. IDI2 catalyzes the 1,3-allylic reorganization of the homoallylic substrate isopentenyl (IPP) to its extremely electrophilic allylic isomer, dimethylallyl diphosphate (DMAPP).

To place an order, please [Click HERE](#).