

NAA10 Human

Description:NAA10 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 255 amino acids (1-235 a.a.) and having a molecular mass of 28.6kDa (the molecular weight on SDS-PAGE will appear higher).NAA10 is fused to a 20 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #:ENPS-165

For research use only.

Synonyms:N-alpha-acetyltransferase 10, N-terminal acetyltransferase complex ARD1 subunit homolog A, NatA catalytic subunit, NAA10, ARD1, ARD1A, TE2, NATD, DXS707.

Source:Escherichia Coli.

Physical Appearance:Sterile Filtered colorless solution.

Amino Acid Sequence:MGSSHHHHHH SSGLVPRGSH MNIRNARPED LMNMQHCNLL
CLPENYQMKY YFYHGLSWPQ LSYIAEDENG KIVGYVLAKM EEDPDDVPHG HITS LAVKRS
HRRGLAQKL MDQASRAMIE NFNAKYVSLH VRKSNRAALH LYSNTLNFQI SEVEPKYYAD
GEDAYAMKRD LTQMADELRR HLELKEKGRH VVLGAIENKV ESKGNSPPSS GEACREEKGL
AAEDSGGDSK DL

Purity:Greater than 90.0% as determined by SDS-PAGE.

Formulation:

NAA10 protein solution (1mg/ml) containing 20mM Tris-HCl buffer (pH 8.0), 5mM DTT, 10% glycerol and 200mM NaCl.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

NAA10 is a member of the acetyltransferase family. NAA10 interacts with NAA15, HIF-1 and with the ribosome. In its binding to HIF-1, NAA10 functions as a protein acetyltransferase by regulating its stability. In various cell lines, NAA10 is downregulated in response to hypoxia. NAA10 is expressed during the course of the development of the brain.

To place an order, please [Click HERE](#).