www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

# **GLRX2** Yeast

Description: Glutaredoxin-2 Saccharamyces cerevisiae Recombinant containing 6x His tag at C-terminus produced in E.Coli is a single, non-glycosylated, Polypeptide chain having a molecular mass of 17 kDa.

Catalog #:ENPS-367

For research use only.

Synonyms: Thioltransferase, Glutathione-dependent oxidoreductase 2, TTR, TTR1, GLRX2, GRX2, GRX-2, GLRX-2, Glutaredoxin 2.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered clear colorless solution.

Purity: Purity of GRX2 is greater than 90% as determined by SDS-PAGE.

#### Formulation:

Glutaredoxin-2 solution contains 25mM Tris-HCl pH-7.5 & 0.01% Na Azide.

## Stability:

1 week at 2-10°C. For long term store at -20 to -80°C.

## Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drµgs, agricultural or pesticidal products, food additives or household chemicals.

## Introduction:

GLRX2 is a multifunctional enzyme with glutathione-dependent oxidoreductase, glutathione peroxidase and glutathione S-transferase (GST) activity. The disulfide bond functions as an electron carrier in the glutathione-dependent synthesis of deoxyribonucleotides by the enzyme ribonucleotide reductase. In addition, it is also involved in reducing cytosolic protein- and non-protein-disulfides in a coupled system with glutathione reductase. Required for resistance to reactive oxygen species (ROS) by directly reducing hydroperoxides and for the detoxification of ROS-mediated damage. Glutaredoxins are a family of glutathione-dependent hydrogen donors that participate in a variety of cellular redox reactions.

To place an order, please Click HERE.





