

MGMT Human

Description: MGMT Human Recombinant fused to 20 amino acid His Tag at N-terminal produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 227 amino acids (1-207) and having a molecular mass of 23.8 kDa. The MGMT is purified by proprietary chromatographic techniques.

Catalog #:ENPS-396

For research use only.

Synonyms: Methylated-DNA--protein-cysteine methyltransferase, 6-O-methylguanine-DNA methyltransferase, O-6-methylguanine-DNA-alkyltransferase, MGMT.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered clear colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MDKDCEMKRT TLDSPLGKLE LSGCEQGLHE IKLLGKGTS AADAEVPAPA AVLGGPEPLM QCTAWLNAYF HQPEAIEFP VPAFHHPVFQ QESFTRQVLW KLLKVVKGFE VISYQQLAAL AGNPKAARAV GGAMRGNPVP ILIPCHRVC SSGAVGNYS GLAVKEWLLA HEGHRLGKPG LGGSSGLAGA WLKGAGATSG SPPAGR N.

Purity: Greater than 95.0% as determined by SDS-PAGE.

Formulation:

The MGMT solution contains 20mM Tris-HCl pH-7.5, 1mM DTT and 10% glycerol.

Stability:

MGMT Recombinant Human although stable at 4°C for 30 days, should be stored desiccated below -20°C for periods greater than 30 days. Please avoid freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

MGMT is an enzyme that repairs O-6-methylguanine, a mutagenic DNA base damaged by endogenous and environmental alkylating agents and takes part in the cellular defense against the biological effects of O-6-methylguanine in DNA. MGMT repairs alkylated guanine in DNA by stoichiometrically transferring the alkyl group at the O-6 position to a cysteine residue in the enzyme. abnormal MGMT expression correlates with the prognosis in human solid cancers. MGMT decrease of expression is correlated with methylation. The human MGMT is a negative regulator of estrogen receptor-mediated transcription upon alkylation DNA damage. MGMT promoter hypermethylation plays an important role in the early steps of colorectal carcinogenesis. Abnormal promoter hypermethylation of MGMT gene is associated with oral squamous cell carcinomas.

To place an order, please Click [HERE](#).