

CDO1 Human

Description: CDO1 Human Recombinant fused with 37 amino acid His tag at N-terminus produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 207 amino acids (1-170 a.a.) and having a molecular mass of 23.9kDa. The CDO1 is purified by proprietary chromatographic techniques.

Catalog #: ENPS-456

For research use only.

Synonyms: Cysteine dioxygenase type 1, Cysteine dioxygenase type I, CDO-I, CDO, CDO1.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MRGSHHHHHH GMASMTGGQQ MGRDLYDDDD KDRWGSHMEQ
TEVLKPRTLA DLIRILHQLF AGDEVNVEEV QAIMEAYESD PTEWAMYAKF DQYRYTRNLV
DQGNGKFNLM ILCWGEHGHS SIHDTNSHC FLKMLQGNLK ETLFAWPDKK SNEMVKKSER
VLRENQCAYI NDSVGLHRVE NISHTEPAVS LHLSPPFDT CHAFDQR.

Purity: Greater than 90.0% as determined by SDS-PAGE.

Formulation:

The CDO1 solution contains 20mM Tris buffer (pH 8.0), 10% glycerol and 1mM DTT.

Stability:

CDO1 although stable 4°C for 4 weeks, should be stored desiccated below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

CDO1 (Cysteine dioxygenase) is a mammalian non-heme iron enzyme that initiates a number of significant metabolic pathways associated with pyruvate and several sulfurate compounds including sulfate, hypotaurine and taurine. CDO1 catalyzes the conversion of L-cysteine to cysteine sulfinic acid (cysteine sulfinate) by incorporation of dioxygen. CDO1 is a vital regulator of cellular cysteine concentrations and has an essential role in maintaining the hepatic concentration of intracellular free cysteine within a proper narrow range. CDO1 is able to alter intracellular cysteine levels and glutathione levels. CDO1 is highly expressed in the liver and placenta. On the other hand CDO1 has a low expression in heart, brain and pancreas. CDO1 can also be detected in hepatoblastoma HepG2 cells.

To place an order, please [Click HERE](#).