

HIV-2 gp-36 (390-702 a.a.)

Description:HIV-2 gp36 34 kDa recombinant- contains the sequence of HIV-2 envelope immunodominant regions gp36, amino acids 390-702. The protein is fused to beta-galactosidase (114 kDa) at N-terminus.

Source:Escherichia Coli.

Physical Appearance:Sterile filtered colorless clear solution.

Amino Acid Sequence:

EQTMVQDDPSTCRGEFLYCNMTWFLNWIENKTHRNYAPCHIKQIINTWHKVGRNVYLPREGEL
SCNSTVTSIIANIDWQNNNQNTNITFSAEVAELRYLELGDYKLVETPIGFAPTKEKRYSSAHGRHTR
GVFVLGFLGFLATAGSAMGAASLTVSAQSRTLLAGIVQQQQQLLDVVKRQQELLRLTVWGTKNL
QARVTAIEKYLQDQARLNSWGCAFRQVCHTTVPWVNDLAPDWDNMTWQEWKQVRYLEA

Purity:Greater than 95.0% as determined by HPLC analysis and SDS-PAGE.

Specificity:

Immunoreactive with all sera of HIV-2 infected individuals.

Formulation:

0.01M Na₂CO₃, 0.01M Na₃EDTA, 0.014 M-mercaptoethanol, 0.02% Sarcosyl.

Stability:

Protein should be stored at 4°C.Refrigerate Upon arrival. DO NOT FREEZE.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Applications:

HIV-2 gp-36 antigen in ELISA and Western blots, excellent antigen for early detection of HIV seroconvertors with minimal specificity problems.

Introduction:

HIV-1 and HIV-2 appear to package their RNA differently. HIV-1 binds to any appropriate RNA whereas HIV-2 preferentially binds to mRNA which creates the Gag protein itself. This means that HIV-1 is better able to mutate. HIV-2 is transmitted in the same ways as HIV-1: Through exposure to bodily fluids such as blood, semen, tears and vaginal fluids. Immunodeficiency develops more slowly with HIV-2.HIV-2 is less infectious in the early stages of the virus than with HIV-1.The infectiousness of HIV-2 increases as the virus progresses.Major differences include reduced pathogenicity of HIV-2 relative to HIV-1, enhanced immune control of HIV-2 infection and often some degree of CD4-independence. Despite considerable sequence and phenotypic differences between HIV-1 and 2 envelopes, structurally they are quite similar. Both membrane-anchored proteins eventually form the 6-helix bundles from the N-terminal and C-terminal regions of the ectodomain, which is common to many viral and cellular fusion proteins and which seems to drive fusion. HIV-1 gp41 helical regions can form more stable 6-helix bundles than HIV-2 gp41 helical regions however HIV-2 fusion occurs at a lower threshold temperature (25°C), does not require Ca²⁺ in the medium, is insensitive to treatment of target cells with cytochalasin B, and is not

www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom
affected by target membrane glycosphingolipid composition.



To place an order, please [Click HERE](#).

Catalog #:HIPS-146

For research use only.

© 2015 NeoScientific



neobiolab | [Term](#) | [Quote/Order](#) | [Services](#) | [Products](#) | [Support](#) | [Corporate](#) | [Contact Us](#)



Copyrights @ 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)

Request a
CATALOG
[CLICK HERE](#)