

HIV-1 gp120 CM

Description: HIV-1 gp120 CM Recombinant- is the external envelope protein, full-length 120 kDa, derived from the env. gene of HIV-1 and glycosylated with N-linked sugars and produced using baculovirus vectors in insect cells. The gp120CM stands for Chaing Mai from Clade E and is M-tropic meaning it binds to CCR5. Purified under conditions that maintain the tertiary structure of the molecule.

Source: Baculovirus, Insect Cells.

Physical Appearance: Sterile filtered colorless clear solution.

Purity: Greater than 90.0% as determined by HPLC analysis and SDS-PAGE.

Formulation:

(311)

Stability:

Recombinant HIV -1 gp120 CM although stable at 4°C for 3 weeks, should be stored below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please avoid freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Human immunodeficiency virus (HIV) is a retrovirus that can lead to a condition in which the immune system begins to fail, leading to opportunistic infections. HIV primarily infects vital cells in the human immune system such as helper T cells (specifically CD4+ T cells), macrophages and dendritic cells. HIV infection leads to low levels of CD4+ T cells through three main mechanisms: firstly, direct viral killing of infected cells; secondly, increased rates of apoptosis in infected cells; and thirdly, killing of infected CD4+ T cells by CD8 cytotoxic lymphocytes that recognize infected cells. When CD4+ T cell numbers decline below a critical level, cell-mediated immunity is lost, and the body becomes progressively more susceptible to opportunistic infections. HIV was classified as a member of the genus Lentivirus, part of the family of Retroviridae. Lentiviruses have many common morphologies and biological properties. Many species are infected by lentiviruses, which are characteristically responsible for long-duration illnesses with a long incubation period. Lentiviruses are transmitted as single-stranded, positive-sense, enveloped RNA viruses. Upon entry of the target cell, the viral RNA genome is converted to double-stranded DNA by a virally encoded reverse transcriptase that is present in the virus particle. This viral DNA is then integrated into the cellular DNA by a virally encoded integrase so that the genome can be transcribed. Once the virus has infected the cell, two pathways are possible: either the virus becomes latent and the infected cell continues to function, or the virus becomes active and replicates, and a large number of virus particles are liberated that can then infect other cells.

References:

Title: Slit2/Robo4 Signaling Modulates HIV-1 gp120-Induced Lymphatic Hyperpermeability. Publication: Zhang X, Yu J, Kuzontkoski PM, Zhu W, Li DY, et al. (2012) Slit2/Robo4 Signaling Modulates HIV-1 gp120-Induced Lymphatic Hyperpermeability. PLoS

www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom
Pathog 8(1): e1002461.

doi:10.1371/journal.ppat.1002461 Link: <http://www.plospathogens.org/article/info%3Adoi%2F10.1371%2Fjournal.ppat.1002461>



Catalog #:HIPS-161

To place an order, please [Click HERE](#).

For research use only.

© 2015 NeoScientific

