www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

# EDN2 Human, His

Description: EDN2 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 177 amino acids (25-178 a.a.) and having a molecular mass of 19.9kDa. EDN2 is fused to a 23 amino acid His-tag at N-terminus & Dry purified by proprietary chromatographic techniques.

Catalog #:HOPS-013

For research use only.

Synonyms: ET2, PPET2, Endothelin-2, ET-2, Preproendothelin-2, EDN2.

Source: Escherichia Coli.

Physical Appearance: Sterile filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MGSQAAATLE QPASSSHAQG THLRLRRCSC SSWLDKECVY FCHLDIIWVN TPEOTAPYGL GNPPRRRRRS LPRRCQCSSA RDPACATFCL RRPWTEAGAV PSRKSPADVF QTGKTGATTG ELLQRLRDIS TVKSLFAKRQ QEAMREPRST HSRWRKR.

Purity: Greater than 85.0% as determined by SDS-PAGE.

#### Formulation:

EDN2 protein solution (1mg/ml) containing 20mM Tris-HCl buffer (pH 8.0), 0.4M urea and 10% glycerol.

# Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

## Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

### Introduction:

Endothelin-2 is a Hypoxia-induced Autocrine Survival Factor for Breast Tumor Cells. The synthesis of EDN2 by human kidney carcinoma cells is decreased by EGF. EDN2 is a chemoattractant for macrophages and THP-1 monocytic cells. Chemotaxis towards EDN2 is via the MAPK pathway: p44 and p42 are phosphorylated when THP-1 cells are stimulated with EDN2. Migration to EDN2 is inhibited by hypoxia and by pertussis toxin. EDN2 leads to activation of macrophages. EDN2 shares a similar peptide sequence with chemokines and may signal via a similar receptor and MAPK-mediated pathway. Furthermore, EDN2 expression by tumors may modulate the behavior of macrophages such that activated cells accumulate in areas of hypoxia.

To place an order, please Click HERE.





