

DnaK SBD

Description: Recombinant DnaK Substrate Binding domain produced in E.Coli is a single, non-glycosylated polypeptide chain containing 132 amino acids and having a molecular mass of 14.6 kDa.

Catalog #: HYP5-015

Synonyms: HSP-70, HSP70, DnaK, Chaperone protein dnaK, Heat shock protein 70, Heat shock 70 kDa protein, groP, grpF, seg, b0014, JW0013.

For research use only.

Source: Escherichia Coli.

Physical Appearance: Sterile filtered colorless solution.

Amino Acid Sequence: MNEDEIQKMV RDAEANA EAD RKFEELVQTR NQGDHLLHST
RKQVEEAGDK LPADDKTAIESALTALETAL KGEDKAAIEA KMQELAQVSQ KLMEIAQQQH
AQQQTAGADASANNAKDDDDVVDAEFEEVKDKK.

Purity: Greater than 95.0% as determined by (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

The protein contains 25mM Tris-HCl, pH7.5, 100mM NaCl, 5mM DTT and 10% Glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

DnaK, originally identified for its DNA replication by bacteriophage I in E. coli is the bacterial hsp70 chaperone. This protein is involved in the folding and assembly of newly synthesized polypeptide chains and in preventing the aggregation of stress-denatured proteins. DnaK (residues 508-638) of the substrate binding domain is α -helical and appears to act as a lid covering the substrate binding cleft. DnaK (amino acid 508-638) was purified to apparent homogeneity by using conventional column chromatography techniques. Additional amino acid (Met) is attached at N-terminus.

To place an order, please [Click HERE](#).